

Name	: Mr. DUMMY	Collected	: 11/7/2017 1:18:00PM
Lab No.	: 133434229	Age: 25 Years	Gender: Male
A/c Status	: P	Ref By : Dr. UNKNWON	Report Status : Final
		Received	: 11/7/2017 1:30:36PM
		Reported	: 18/7/2017 3:22:58PM

Test Name	Results	Units	Bio. Ref. Interval
IHC MARKER(S)	RESULT		
CD 21	Non-immunoreactive, score 0 in neoplastic cells		

SLIDE NO : B/ 2222222/17
SPECIMEN : Biopsy for IHC markers.
CLINICAL HISTORY : --
GROSS : Received 1 formalin fixed paraffin embedded block labelled as 'ABCDE'
IMPRESSION : **Features are consistent with XYZ**

INTERPRETATION

RESULT	SCORE
Non immunoreactive	0
Immunoreactive in 1-25 % cells	1+
Immunoreactive in 26-50% cells	2+
Immunoreactive in 51-75% cells	3+
Immunoreactive in 76-100% cells	4+

COMMENTS

CD21:The B2 antigen, also known as CD21 and CR2, is a 140 kD glycosylated Type 1 integral membrane protein. The CD21 molecule is expressed on mature B lymphocytes, follicular dendritic reticulum cells (FDC), pharyngeal epithelial cells, and possibly on a subset of normal thymocytes. This antigen is also expressed by B lymphocytes in patients with B cell lymphomas, most B cell chronic lymphocytic leukemia (CLL), and a small portion of non-T cell acute lymphoblastic leukemias (ALL).

NOTE

1. Detection system used is Polymer HRP
2. The impression is based on the material submitted and is not a complete surgical pathology report.
3. False negative IHC results due to inadequate fixation of the material sent for evaluation cannot be excluded.

FIXATION REQUIREMENTS

- The volume of formalin fixative should be atleast 10 times the volume of the specimen.
- Decalcification solutions with strong acids should not be used.



LPL - LPL-ROHINI (NATIONAL REFERENCE
LAB)
SECTOR - 18, BLOCK -E ROHINI
DELHI 110085

Name	: Mr. DUMMY	Collected	: 11/7/2017 1:18:00PM
Lab No.	: 133434229	Age: 25 Years	Gender: Male
A/c Status	: P	Ref By : Dr. UNKNWON	Report Status : Final
		Received	: 11/7/2017 1:30:36PM
		Reported	: 18/7/2017 3:22:58PM

Test Name	Results	Units	Bio. Ref. Interval
	<ul style="list-style-type: none">Specimens should be immersed in fixative within 1 hour of the biopsy/resection procedure (time of removal & time of immersion to be mentioned).In all resection (large) specimens, the tumour must be bisected prior to immersion in fixative		

DR. HEMA MALINI AIYER
MD (PATH)
HOD-HISTOPATH/CYTOPATH

-----End of report-----

