

<b>PHOSPHATIDYL SERINE ANTIBODY, IgM, SERUM ( EIA)</b>	<b>Index</b>	<b>&lt; 0.80</b>
--	--------------	------------------

**2006 International Consensus statement on Classification of definite APS**

<b>Clinical Criteria</b>	<b>Laboratory Criteria</b>
Arterial / Venous thrombosis	Cardiolipin antibodies ( aCL)
Fetal Loss	Beta 2 Glycoprotein 1 Antibodies
Premature birth	Lupus Anticoagulant ( LA)

**Note:** APS is established if at least 1 Laboratory criteria and 1 Clinical criteria are met. The Laboratory criteria should be present on two or more occasions 12 weeks apart for diagnosing APS.

**Comments**

Phospholipid antibody is a quantitative assay to screen the presence of autoantibodies against cardiolipin, phosphatidyl serine, phosphatidyl inositol & phosphatidic acid in the diagnosis of an increased risk of thrombosis in patients with Systemic lupus erythematosus (20-35%) and other lupus like disorders.

This test is also used to diagnose Anti-phospholipid syndrome in patients with recent miscarriage ( 11-22% in all trimesters), Pulmonary hypertension, Non-vegetative endocarditis, Livedoreticularis, Stroke at young age and Deep vein thrombosis. Measurement of Phosphatidylserine antibodies in addition to Antiphospholipids enables a more specific diagnosis.