

Name	: DUMMY	Collected	: 29/8/2017 12:00:00AM
Lab No.	: 135091659	Age: 45 Years	Gender: Male
A/c Status	: P	Ref By : Dr. UNKNWON	Report Status : Final
		Received	: 29/8/2017 10:02:20AM
		Reported	: 29/8/2017 10:55:35AM

Test Name	Results	Units	Bio. Ref. Interval
<b>CARDIOLIPIN ANTIBODIES PANEL</b>			
CARDIOLIPIN ANTIBODY, IgA ,SERUM (EIA)	4.00	APL	<12.00

**Interpretation**

RESULT IN APL	REMARKS
<12	Negative
12-20	Equivocal
20-80	Low Positive
>80	High Positive

**Comments**

Anticardiolipin antibodies(ACA) belong to the group of Antiphospholipid antibodies which are positive in 30-40% cases of Systemic lupus erythematosus and also in patients with other Rheumatic diseases. Presence of cardiolipin antibodies is considered to be of significant diagnostic relevance in cases of Venous/Arterial thrombosis, Thrombocytopenia, Livedo reticularis, Habitual abortions and Neurological manifestations. Elevated ACA levels are also seen in patients with Cardiovascular insufficiency and Myocardial infarction.

CARDIOLIPIN ANTIBODY, IgG, SERUM (EIA)	5.00	GPL	<15.00
--	------	-----	--------

**Interpretation**

RESULT IN GPL	REMARKS
<15	Negative
15-20	Equivocal
20-80	Low Positive
>80	High Positive

**Comments**

Anticardiolipin antibodies(ACA) belong to the group of Antiphospholipid antibodies which are positive in 30-40% cases of Systemic lupus erythematosus and also in patients with other Rheumatic diseases. Presence of cardiolipin antibodies is considered to be of significant diagnostic relevance in cases of Venous/Arterial thrombosis, Thrombocytopenia, Livedo reticularis, Habitual abortions and Neurological manifestations. Elevated ACA levels are also seen in patients with Cardiovascular insufficiency and



Name	: DUMMY	Collected	: 29/8/2017 12:00:00AM
Lab No.	: 135091659	Age: 45 Years	Gender: Male
A/c Status	: P	Ref By : Dr. UNKNWON	Report Status : Final
		Received	: 29/8/2017 10:02:20AM
		Reported	: 29/8/2017 10:55:35AM

**Test Name** **Results** **Units** **Bio. Ref. Interval**  
 Myocardial infarction. Results must be correlated with the history and clinical findings of the patient.

<b>CARDIOLIPIN ANTIBODY, IgM, SERUM (EIA)</b>	12.00	MPL	<12.50
---	-------	-----	--------

**Interpretation**

RESULT IN MPL	REMARKS
<12.50	Negative
12.50-20	Equivocal
20-80	Low Positive
>80	High Positive

**Comments**

Anticardiolipin antibodies (ACA) belong to the group of Antiphospholipid antibodies which are positive in 30-40% cases of Systemic lupus erythematosus and also in patients with other Rheumatic diseases. Presence of cardiolipin antibodies is considered to be of significant diagnostic relevance in cases of Venous/Arterial thrombosis, Thrombocytopenia, Livedo reticularis, Habitual abortions and Neurological manifestations. Elevated ACA levels are also seen in patients with Cardiovascular insufficiency and Myocardial infarction.

*Ritu*

Dr. Ritu Nayar  
 MD (Microbiology)  
 Deputy HOD Microbiology & Serology

*Shalabh Malik*

Dr. Shalabh Malik  
 MD (Microbiology)  
 National Head - Microbiology & Serology

-----End of report -----

